

BIROn - Birkbeck Institutional Research Online

Engels, Stefan and Medeiros, A.S. and Axford, Y. and Brooks, S.J. and Heiri, O. and Luoto, T.P. and Nazarova, L. and Porinchu, D.F. and Quinlan, R. and Self, A.E. (2020) Temperature change as a driver of spatial patterns and long-term trends in chironomid (Insecta: Diptera) diversity. *Global Change Biology* 26 (3), pp. 1155-1169. ISSN 1354-1013.

Downloaded from: <https://eprints.bbk.ac.uk/id/eprint/29416/>

Usage Guidelines:

Please refer to usage guidelines at <https://eprints.bbk.ac.uk/policies.html>
contact lib-eprints@bbk.ac.uk.

or alternatively

Temperature change as a driver of spatial patterns and long-term trends in chironomid

(Insecta: Diptera) diversity

Running head: Temperature change and chironomid diversity

List of authors: Stefan Engels^{1*}, Andrew S. Medeiros², Yarrow Axford³, Stephen J. Brooks⁴, Oliver Heiri⁵, Tomi P. Luoto⁶, Larisa Nazarova^{7,8,9}, David F. Porinchu¹⁰, Roberto Quinlan¹¹, Angela E. Self⁴

List of institutes:

¹ Department of Geography, Birkbeck University of London, London WC1E 7HX, UK; s.engels@bbk.ac.uk

² School for Resource and Environmental Studies, Dalhousie University, 6100 University Ave, Halifax, NS B3H 4R2, Canada; andrew.medeiros@dal.ca

³ Northwestern University, Department of Earth and Planetary Sciences, 2145 Sheridan Road, Evanston IL USA 60208; axford@northwestern.edu

⁴ Department of Life Sciences, Natural History Museum, Cromwell Road, London SW7 5BD, UK; S.Brooks@nhm.ac.uk; angela.e.self@btinternet.com

⁵ Geoecology, Department of Environmental Sciences, University of Basel, Klingelbergstrasse 27, CH-4056 Basel, Switzerland; oliver.heiri@unibas.ch

⁶ Faculty of Biological and Environmental Sciences, Ecosystems and Environment Research Programme, University of Helsinki, Niemenkatu 73, FI-15140 Lahti, Finland; tomi.luoto@helsinki.fi

⁷ Institute of Earth and Environmental Science, University of Potsdam, Karl-Liebknecht-Str. 24-25, 14476 Potsdam-Golm, Germany; larisa.nazarova@uni-potsdam.de

⁸ Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research, Research Unit Potsdam, Telegrafenberg A43, 14473 Potsdam, Germany

⁹ Kazan Federal University, Kremlyovskaya str. 18, 420018 Kazan, Russia

¹⁰ Department of Geography, University of Georgia, Athens GA 30602, USA; porinchu@uga.edu

¹¹ Department of Biology, York University, 4700 Keele St, Toronto, ON M3J 1P3, Canada; rquinlan@yorku.ca

* Author for correspondence: Stefan Engels; tel: 0044 (0)20 30738457; s.engels@bbk.ac.uk

Key words: biodiversity, climate warming, palaeoecology, Quaternary, Arctic, freshwater ecosystems, insects

Paper type: Primary Research Article

Abstract

Anthropogenic activities have led to a global decline in biodiversity, and monitoring studies indicate that both insect communities and wetland ecosystems are particularly affected. However, there is a need for long-term data (over centennial- or millennial timescales) to better understand natural community dynamics and the processes that govern the observed trends. Chironomids (Insecta: Diptera: Chironomidae) are often the most abundant insects in lake ecosystems, sensitive to environmental change, and, because their larval exoskeleton head capsules preserve well in lake sediments, they provide a unique record of insect community dynamics through time. Here, we provide the results of a meta-data analysis of chironomid diversity across a range of spatial and temporal scales. First, we analyse spatial trends in chironomid diversity using Northern Hemispheric datasets overall consisting of 837 lakes. Our results indicate that in most of our datasets summer temperature (T_{jul}) is strongly associated with spatial trends in modern-day chironomid diversity. We observe a strong increase in chironomid alpha diversity with increasing T_{jul} in regions with present day T_{jul} between 2.5-14 °C. In some areas with $T_{\text{jul}} > 14$ °C chironomid diversity stabilises or declines. Second, we demonstrate that the direction and amplitude of change in alpha diversity in a compilation of subfossil chironomid records spanning the last glacial-interglacial transition (~15,000-11,000 years ago) are similar to those observed in our modern data. A compilation of Holocene records shows that during phases when the amplitude of temperature change was small, site-specific factors had a greater influence on the chironomid fauna obscuring the chironomid diversity-temperature relationship. Our results imply expected overall chironomid diversity increases in colder regions such as the Arctic under sustained global warming, but with complex and not necessarily predictable responses for individual sites.

Introduction

Key environmental parameters such as greenhouse gas concentrations, ocean acidity, and erosion rates are now beyond their previous Holocene range of variability as a result of human activity (Corlett, 2015; Steffen et al., 2015). These environmental stressors are leading to dramatic changes in the structure and composition of biological communities across local to global spatial scales and to a decrease in global biodiversity (e.g. Ripple et al., 2017). For example, the invertebrate fauna of northwest Europe is showing signs of a dramatic decrease in abundance and diversity (Hallmann et al., 2017; Bidau, 2018; Jourdan et al., 2018; Leather 2018; Rada et al., 2019) and freshwater ecosystems are showing a particularly high rate of biodiversity loss (Dudgeon et al., 2006).

Most records that are used to assess the effects of anthropogenic drivers (e.g. climate variability, eutrophication, toxins, impacts associated with invasive species) on biodiversity are based on decadal-scale ecological data (Vellend et al., 2013; Birks et al., 2016a; W.W.F., 2016). In many natural settings it has been difficult to disentangle natural variability from the effects of human-induced ecosystem change (Loreau et al., 2001; Willis & Birks, 2006) as a result of these short time-scales typically used. Datasets that cover much longer timescales, e.g. several centuries to millennia, can help to provide a broader temporal context for current and future trends in biodiversity and facilitate the development of effective conservation strategies (Willis & Birks, 2006; Gregory-Eaves & Beisner, 2011; Birks et al., 2016a). Willis & Birks (2006) and Birks et al. (2016a) provide examples of the role that palaeoecological datasets can play in providing such centennial- to millennial-scale time series, and how these data can provide new insights into the complex relationships that exist between biodiversity, ecosystem functioning, and environmental variability. Despite the fact that many natural archives are available for study by palaeo- and neo-ecologists, these sources of information have rarely been used to infer past changes in biodiversity within the context of human

impact and the ‘Anthropocene’ (Gregory-Eaves & Beisner, 2011; Corlett, 2015). We therefore have a limited understanding of long-term trends (10^2 - 10^5 years) in biodiversity, including insect diversity, and, as a result, it remains difficult to disentangle effects of anthropogenic disturbances from natural trends and internal variability.

Palaeoecological records describe the development of ecosystems over timescales ranging from a few decades to several hundreds of thousands of years (e.g. Smol et al., 2005). They are typically derived from lake deposits, peat bogs, marine sediments, or other natural archives. Fossils encountered in these materials can be used to reconstruct past ecological and environmental change and provide estimates of past biodiversity for different parts of the ecosystem (e.g. terrestrial vegetation, algal flora, invertebrate fauna) (e.g. Smol et al., 2005).

A method that has provided insights into ecological, environmental and climatic development across the last glacial/interglacial cycle is the analysis of subfossil chironomid remains preserved in lake sediments (e.g. Brooks, 2006; Walker & Cwynar, 2006). Chironomids (Insecta: Diptera: Chironomidae) are holometabolous insects, and the larval stage of most species is aquatic. In lakes, chironomids are usually the most abundant macroinvertebrate group, and they are a critical component of the benthic community, contributing to processes essential for lake ecosystem functioning (Hölker et al., 2015), including biogeochemical cycling of carbon and nutrients. Parts of their larval exoskeleton (the head capsules) are well-preserved in lake sediments (e.g. Brooks et al., 2007). Subfossil chironomid head capsules are abundant in lake sediments, often in concentrations of 50-100 head capsules per gram of sediment, can be identified to genus, species morphotype, or species level, and provide a record of changes in the chironomid fauna of a lake through time. By comparing subfossil chironomid assemblages to their modern-day distribution patterns, quantitative reconstructions of changes in their past environment can be made. For instance, chironomids have been used to quantitatively reconstruct past changes in temperature

(Medeiros et al., 2012; Heiri et al., 2014; Brooks et al., 2016), water depth (Engels et al., 2012; Nazarova et al., 2017), salinity (Dickson et al., 2014), hypolimnetic oxygen (Quinlan & Smol, 2001), and several other environmental variables. Subfossil chironomids encountered in lake sediment records can additionally be used to track changes in past species richness or evenness of the fauna through time (e.g. Engels et al., 2010). However, palaeoecological datasets describing chironomid faunal dynamics on centennial to millennial timescales have not yet been compiled to produce systematic and super-regional studies of past diversity patterns.

While it is often assumed that the global decline in biodiversity is accompanied by declines at finer spatial scales, local biodiversity has been shown to have been stable in many regions even when global biodiversity trends show significant decreases (Cardinale et al., 2018; Primack et al., 2018). It therefore follows that biodiversity at large spatial scales is not necessarily an additive function of biodiversity at smaller scales (Vellend et al., 2013) and that investigating biodiversity at a range of spatial scales is necessary to understand the ongoing changes in biodiversity as a result of climatic and environmental change (Primack et al., 2018). While palynological records provide long-term trends of regional changes in vegetation (e.g. Birks et al., 2016a), there are currently no studies that assess long-term diversity change on finer spatial scales, such as individual aquatic ecosystems. Therefore, this study aims to contribute toward filling the gap in our understanding of long-term trends in local-scale biodiversity changes and to provide a long-term perspective to the recent decline in insect diversity. Specifically, we aim to assess long-term changes in chironomid diversity, a keystone member of freshwater ecosystems, by:

- (1) Determining the relationship between diversity of (subfossil) chironomid assemblages in sediment samples and environmental parameters, with an emphasis on

the effects of summer temperature on chironomid assemblages across the mid- and high-latitudes of the northern hemisphere (reflecting the availability of datasets); (2) Analysing spatiotemporal changes in chironomid diversity across a range of timescales; and (3) Assessing the suitability of subfossil chironomid assemblages as an approach to reconstruct changes in diversity over longer timescales.

Materials and methods

Materials

We determine, for the first time, spatial trends in chironomid diversity in chironomid-climate calibration datasets from Norway (Brooks & Birks, 2001; Brooks, 2006), Russia (Nazarova et al., 2011; 2015; Self et al., 2011), Canada (Gajewski et al., 2005; Porinchu et al., 2009; Fortin & Gajewski, 2011; Medeiros & Quinlan, 2011), Switzerland (Lotter et al., 1997; Heiri & Lotter, 2010) and Finland (Luoto 2009a; 2009b) (Fig. 1; Supplementary Table S1). Samples presented here from the Canadian dataset have been re-identified where necessary and the taxonomy of the datasets analysed here follows Brooks et al. (2007).

Chironomid-climate calibration datasets document the relationship between chironomids and their contemporaneous environment (Brooks, 2006), often with a specific focus on the influence of summer air temperature on chironomid distribution patterns (e.g. Heiri et al., 2014). As such, these datasets capture a wide (summer) temperature range, whereas the impact of other environmental variables (e.g. lake water pH) is typically minimised. As a result of this sampling approach, the datasets provide excellent resources to study the temperature-chironomid diversity relationship. However, as the datasets cover a range of different secondary gradients (e.g. Medeiros et al., 2015) and include measurements of different environmental variables (see Supplementary Table S1), testing the influence of

environmental parameters other-than-temperature on chironomid diversity will be more empirical.

To determine temporal trends in chironomid diversity we selected northern hemisphere sites with clear information on the taxonomic resolution, and for which we had access to the original count data (Supplementary Table S2, S3). The sites presented in this study cover different timescales, ranging from records that date back as far as 200,000 cal yr BP to records that cover the last few centuries. We specifically focus on the last glacial-interglacial transition (LGIT; ca. 14,500-11,600 calendar years before the present (cal yr BP)), the current interglacial, i.e. the Holocene (ca. 11,600 cal yr BP to the present) and high-resolution records covering the last few centuries. For the selection of the Holocene and recent records used in this study (Supplementary Table S3) we focussed on sites that are relatively removed from local human impacts.

The LGIT was characterised by a number of climate events in the North Atlantic region. The general trend of climate warming was interrupted by a major cold phase (stadial) known as the Younger Dryas (YD), which occurred between ca. 12,650 and 11,600 cal yr BP (e.g. Litt et al., 2001). The relatively warm Bølling-Allerød interstadial (B/A; ca 14,500-12,650 cal yr BP) preceding the YD was punctuated by at least two decadal to centennial scale climate cooling events, the Gerzensee Oscillation (GZO or Greenland Interstadial event (GI)-1b) and the Older Dryas (or GI-1d; Lotter et al., 1992). The YD has been palynologically identified in each of the LGIT records presented in this paper and several of the records also identified the Gerzensee Oscillation and the Older Dryas (depending on location of the sites, temporal sampling resolution and proxy sensitivity). Supplementary Table S2 provides more details on the sites used in this study.

Numerical analyses

We focus on two biodiversity measures that are relevant to subfossil chironomid assemblages: (a) alpha diversity, here defined as the taxon richness of an assemblage at one point in time and space, and (b) compositional turnover, i.e. changes in assemblage composition across time (cf. McGill et al., 2015).

First, we analysed spatial trends in alpha diversity by calculating taxon richness using rarefaction analysis for each of our modern datasets. Rarefaction simulates a random selection without replacement, estimating the taxon richness for each sample using a fixed count size, which typically equals the smallest count sum recorded in a sequence of samples (e.g. Birks & Line, 1992). Results of rarefaction analyses can be quantitatively compared between samples and datasets with different counts. Prior to rarefaction analysis, we rounded the counts in our datasets to integers and set our cut-value to a count sum of 40 head capsules for all analyses. Whilst chironomid studies often report a minimum count sum of 50 head capsules per sample, we noted that some of our fossil sites had depth-intervals where this number was not consistently reached. Instead of amalgamating or removing these samples and thus losing information, we decided to set a cut-value of 40 head capsules in order to retain more samples. Test runs comparing results of changing cut-values from 50 to 40 head capsules showed no substantial differences in the results and did not change interpretations of patterns and trends. Sites that had count values below 40 head capsules were removed prior to analysis. Rarefaction analysis was implemented using the vegan package in R (Oksanen et al., 2015).

Visual inspection of scatterplots of selected environmental variables against alpha diversity revealed the existence of non-linear but monotonic relationships. We subsequently calculated Spearman rank-order correlation coefficients between the rarefaction values and

environmental observations for each modern dataset (Table 1). The results indicated that generally there was a significant positive association between July air temperature (T_{Jul}) and alpha diversity. We produced scatter plots of chironomid diversity (estimated using rarefaction analysis) as a function of July air temperature and used Generalized Additive Modelling (GAM) to summarise trends in the data (Simpson, 2018) (see Supplementary Information for R code).

Second, we reconstructed trends in alpha diversity as well as in compositional turnover (as an approximation for temporal beta diversity) for each of our fossil records. Prior to rarefaction analysis, samples with < 40 head capsules were amalgamated with adjacent samples, making sure that samples were only combined within lithological units. If samples could not be combined with adjacent samples they were left out of the analysis.

While beta diversity is a commonly used component in (palaeoecological) biodiversity studies, there is no consensus on how to derive and interpret beta diversity (e.g. Jost 2007; Marcon et al., 2012; Birks et al., 2016b). Beta diversity, or the change in species composition per unit space or time (McGill et al., 2015), is often quantified as a rate-of-change or a turnover metric (Birks, 2012; Birks et al., 2016a). Even though estimating assemblage compositional turnover along a temporal gradient does not provide a true diversity component but rather a measure of compositional complexity (Tuomisto, 2010; Birks et al., 2016b), compositional turnover is commonly interpreted to approximate temporal changes in beta diversity, and compositional turnover techniques have become popular tools in palaeoecological studies (e.g. Smol et al., 2005; Birks & Birks, 2008; Colombaroli & Tinner, 2013; Feurdan et al., 2013; Stivrins et al., 2016). The difference in scores on a Detrended Canonical Correspondence Analysis (DCCA) axis between subsequent samples

can be used to highlight times of increased turnover, reflecting increased beta diversity between samples.

In this study we follow the examples of Smol et al. (2005) and Birks & Birks (2008) and estimate changes in beta diversity through calculating DCCA axis 1-scores on a temporal gradient. DCCA results are robust to differences in sample numbers for comparable time periods (Birks, 2007), as is the case for e.g. our LGIT records. Changes in compositional turnover were determined by running DCCAs for each of our fossil sites with time as the constraining variable, or, when no robust chronology was available, using sample depth as the constraining variable. In accordance with previously published results, the percent-abundance datasets were square-root transformed prior to DCCA, and we applied detrending by segments with no down-weighting of rare taxa and non-linear rescaling in our analyses. DCCAs were performed using CANOCO v4.5 (ter Braak and Šmilauer, 2002).

Finally, we quantitatively compare the rarefaction results for the Norwegian calibration dataset to the LGIT reconstructions to assess whether the effects of temperature on alpha diversity, as observed in modern-day chironomid assemblages, are comparable to those of the distant past. To do so, we assessed the change in rarefaction (ΔRF) with changing T_{jul} (ΔT_{jul}) across the entire temperature interval included in the Norwegian calibration dataset. In a first step, we calculated the average RF (RF_{ave}) value for each consecutive 1-°C bin (e.g. 3.00-3.99 °C ($T_{3.5C}$)) in the Norwegian calibration dataset. We then calculated changes in RF (ΔRF) for each possible combination of 1-°C bins across the 3-16 °C temperature interval covered by the Norwegian calibration dataset. This approach allows us to observe whether e.g. a change of 2 °C has the same effects on the taxonomic diversity of a lake situated e.g. at 8 °C as for a lake situated at e.g. 14 °C, as well as to compare the effects an increase of 2 °C would have compared to the effects of an increase of 4 °C at the same site. We used the

geom_raster function as included in the R ggplot2 package to produce a continuous plot of changes in RF (Δ RF) for each absolute temperature (e.g. $T = 3.5\text{ }^{\circ}\text{C}$) and each possible increase in temperature (ΔT).

We subsequently calculated the change in alpha diversity associated with the two major climate transitions in our LGIT records, the Allerød-YD and the YD-Holocene transitions. We assessed RF_{ave} for each climate zone in each LGIT record, avoiding periods of transition and the effects of short-scale oscillations such as the Older Dryas, and calculated Δ RF values by comparing RF_{ave} for subsequent climate zones (e.g. YD versus Holocene). We use published quantitative LGIT T_{jul} inferences (see Supplementary Table S2) to assess the average temperature during each interval (e.g. T_{YD}) and calculated the change in temperature between subsequent intervals (ΔT) to be able to compare the LGIT results to the modern-day patterns in alpha diversity. Using this approach, we can assess what the change in alpha diversity was at a given starting temperature and under a particular climate forcing (i.e. temperature difference between consecutive time intervals). These inferences for the fossil records can be quantitatively compared to the inferences based on the modern dataset to assess whether fossil and present trends in the chironomid diversity-temperature relationship are of a similar magnitude. We refer to the supplementary information for more details on the procedure.

The T_{jul} records used in this calculation are chironomid-based temperature inferences and are based on the same datasets as the alpha diversity reconstructions, and as such are not independent. However, in the absence of independent temperature estimates this is currently the only method that allows the quantitative comparison of our modern data to our fossil reconstructions.

Results and interpretation

Spatial trends in chironomid diversity

All datasets show similar TR values ranging between minima of 1-5 to maximum values over 20 (Fig. 2). Chironomid alpha diversity generally shows an increasing trend with increasing T_{jul} for four out of the five regional datasets (Fig. 2). In the Norwegian and Canadian datasets the increase in alpha diversity is steepest in the coldest part of the temperature gradient (ca. 3-7 °C interval), after which a more moderate increase in diversity can be seen. For some of the datasets that cover temperatures exceeding ca. 14 °C (e.g. Russia, Switzerland) we observe a stabilising or decreasing trend in alpha diversity with higher temperatures. Three of the datasets show additional variation superimposed on the general trend of increasing diversity with increasing temperature, which might be the result of sampling density along the temperature gradient, site-specific factors resulting in a lower alpha diversity in individual lakes, or it might be a true reflection of decreasing alpha diversity for currently unidentified reasons. The Finnish dataset captures a smaller temperature gradient than the other datasets and GAM shows a highly variable but non-directional trend in alpha diversity. However, as in the other datasets, some of the lakes with mean July air temperature higher than 14 °C are characterized by relatively low alpha diversity. Explanations for the general increase in chironomid diversity with temperature in most of our datasets could include availability of more and higher-quality food sources with increasing temperatures. For instance, the lengthening of the ice-free period in Arctic lakes has been shown to lead to more diverse and ecologically complex algal communities (Griffiths et al., 2017). Additionally, lakes can support more complex and varied habitats with increasing temperatures as e.g. more complex aquatic macrophyte communities, a major structuring element in lake ecosystems, may occur in warmer lakes. Additionally, more pronounced thermal stratification under warmer climates may lead to the parallel occurrence of both summer warm and summer cold habitats in the same lake, whereas this may not be the case in colder climatic conditions. The stabilising or

even decreasing trends in alpha diversity at the warm end of the gradient could be explained by temperature-related effects such as seasonal stratification in combination with oxygen depletion, leading to the elimination of some chironomid species. Similarly, warmer lakes could experience a loss of aquatic macrophyte habitat due to phytoplankton blooms.

Comparisons of alpha diversity against environmental factors reveal that T_{jul} is the factor that is most strongly associated with chironomid diversity in four of our datasets (Table 1), with Spearman rank correlation coefficients between chironomid diversity and T_{jul} reaching a maximum value of $\rho=0.79$. Many of the chironomid datasets used in this study have been specifically designed to capture a large temperature gradient while aiming to minimise the influence of other environmental parameters, potentially partially explaining why water depth, pH or conductivity do not show strong relationships with chironomid diversity.

There is a positive association between TOC/DOC and chironomid diversity in the datasets for which TOC/DOC data is available (ρ ranges between 0.35 and 0.70). However, TOC/DOC is strongly correlated to T_{jul} in e.g. the Swiss and Norwegian datasets (Lotter et al., 1997; 1998; Brooks & Birks, 2001), and it is therefore unclear to what extent variations in temperature and variations in TOC/DOC independently drive spatial patterns in chironomid diversity. Similarly, there is a positive association between TP and chironomid diversity but it is unclear what the unique effects of TP are, independent of temperature. To further test the potential association between chironomid diversity and TP we applied rarefaction analysis to a lake dataset from Ontario (USA) by Quinlan & Smol (2001) which covers a gradient from ultraoligotrophic lakes to eutrophic lakes whilst covering a relatively short temperature gradient. The results indicate that there is no statistically significant association between chironomid diversity and TP ($\rho = 0.072$, $p = 0.573$) in this dataset.

Similarly, by running additional tests on a chironomid dataset from New England, US (Engels & Cwynar, 2011) as well as a dataset of lakes and reservoirs from New Jersey and New York states, US (Wazbinski & Quinlan, 2013) we further explored whether the absence of a strong correlation between water depth and chironomid diversity could be related to dataset design. The distribution of many individual chironomid taxa in the dataset by Engels & Cwynar (2011) is significantly correlated to water depth, likely attributable to factors like habitat differentiation, taphonomical processes, and oxygen availability. Analysis of both datasets shows only a weak positive association between chironomid diversity and water depth of $\rho = 0.08$ ($p=0.37$) for the Engels & Cwynar (2011) dataset, whereas the Wazbinski & Quinlan (2013) dataset shows a negative association of $\rho = -0.38$ ($p=0.01$) between water depth and chironomid diversity. The combined results suggest that unlike summer temperature, water depth is not a strong driver of chironomid diversity at the observed spatial scales, but that more research is needed to further explore the relationship between chironomid diversity and water depth.

Temporal trends in chironomid diversity

Last glacial–interglacial transition (LGIT)

Four of the six UK LGIT records show a distinct change in diversity during the YD (Fig. 3a), either as a decrease in alpha diversity (e.g. Ashik; Fig. 3a) or as high compositional turnover (e.g. Sunbiggin Tarn; Fig. 3a). The records from Abernethy and Urswick Tarn furthermore show a decrease in both alpha diversity and increased compositional turnover during the multi-decadal cold interval associated with the Older Dryas (Fig. 3a). The records from Ashik and Whitrig Bog show high compositional turnover during the Older Dryas, but no variation in alpha diversity. The Older Dryas has not been identified in the Little Hawes Water and Sunbiggin Tarn records, and although there are oscillations in the compositional turnover

records of these sites, it is unclear whether these are associated with Older Dryas climate change or if they are the result of other factors.

The six LGIT sites from the European mainland (Fig. 3b) are located on a N-S transect (Fig. 1). The southernmost two of these sites (Aubrac, Gerzensee) capture different parts of the LGIT interval, but neither site shows abrupt changes in diversity associated with the onset of the YD or the Holocene (Fig. 3). The records from Hijkermeer and Kråkenes show a decrease in alpha diversity as well as increased compositional turnover at the onset of the YD, whereas the diversity records for the sites from northern Europe (Lusvatnet, Jansvatnet) do not show any response to this large-scale climate cooling. A potential explanation for this discrepancy could lie in the fact that the amplitude of climate change showed regional differences across northwest Europe, with the most pronounced temperature change observed for the British Isles (e.g. Heiri et al., 2014), and that the change in temperature associated with the onset of the YD was not large enough to affect chironomid diversity across the European continent. All four central to northern European mainland sites show a distinct change in their alpha diversity and/or in compositional turnover records at the onset of the Holocene. The amplitude of climate change across this interval was large and occurred within a few decades (e.g. Brauer et al., 1999), and the direct effects of temperature most likely strongly influenced the composition of the freshwater invertebrate fauna across large parts of Europe (Heiri et al., 2014).

In summary, most of the LGIT records document decreasing chironomid diversity during colder periods, both during major climate oscillations (YD) as well as during shorter-lived climate oscillations (Older Dryas). The responses are not uniform across all sites, with some sites showing a decrease in alpha diversity and an increase in compositional turnover, but other sites only showing changes in one of the diversity parameters.

Holocene

The combined results for the Holocene records illustrate the complex relationship between chironomid diversity and the environment, where factors other than temperature override the diversity-temperature relationship that is apparent in the LGIT records. For instance, the chironomid diversity reconstructions for three Holocene chironomid records from the Kamchatka Peninsula show major differences between the sites (Fig. 4a). Lifebuoy Lake (Solovieva et al., 2015) shows changes in diversity and compositional turnover in the early Holocene, Pechora Lake (Andrén et al., 2015) shows increasing variability in the latter part of the record, and Olive-backed Lake (Self et al., 2015) shows a decreasing trend in alpha diversity throughout the entire Holocene. We suggest that for the Holocene, when temperature fluctuations for large parts of the Northern Hemisphere are generally assumed to have been smaller and more gradual than those of the LGIT (e.g. Bartlein et al., 2015), factors other than temperature (e.g. vegetation change, soil development) were more important drivers of changes in chironomid diversity at these sites.

Similarly, sites within relatively small regions like Norway and Finland show differences in both alpha diversity and compositional turnover trends during the Holocene, whereas climate change over these regions can be expected to be fairly homogeneous. Again, we attribute these differences in diversity trends to site-specific changes in the environment exerting a stronger influence on the chironomid fauna than climate change. For instance, the amplitude of temperature change during the Holocene across northern Europe was not as great as during the LGIT, and the chironomid fauna of some lakes, e.g. Giltjärnen, most likely responded most strongly to changes in the local environment (e.g. lake water pH, local peat development) rather than to temperature change (Antonsson et al., 2006).

From the North American sites, the Lake Hill record was retrieved from the remote St. Paul Island (Alaska, USA) and is assumed to have been primarily influenced by climate rather than anthropogenic factors (Wooller et al., 2018). It shows a decrease in alpha diversity around 6,000 cal yr BP (Fig. 4d). Similarly, while not reaching as far back in time as the Lake Hill record, the TK2 site from the Central Canadian Arctic shows high alpha diversity in the early part of the Holocene, with a decrease in alpha diversity and higher compositional turnover in the middle Holocene (ca 7,000 cal yr BP) (Porinchu et al., 2019). We suggest that these declines in diversity are related to regional climate cooling (Kaufman et al., 2004; Clegg et al., 2011).

Recent trends

The complexity of the chironomid diversity-temperature relationship is also highlighted by recent records from northern North America (Fig. 5). While diversity increases with periods of recent anthropogenic warming in Baker Lake (BL01) and Lake T02, it decreases in Lake T15 (Fig. 5). The thermokarst basin of Lake T15 was shown to expand during a period that predated the warming trends for the southern Seward Peninsula in Alaska (Medeiros et al., 2014). The expansion of the thermokarst basin likely altered available habitat, including macrophyte growth and distribution, and as a result the chironomid assemblage shifted from a diverse fauna to a fauna dominated by a single taxon (Medeiros et al., 2014).

Deep-time (10^5 years) records

The lacustrine record from lake Billjakh (east Siberia, Russia; Diekmann et al., 2016) covers the time interval from the middle of the last glacial (MIS-3) to the present. The alpha diversity record of Billjakh shows highest diversity in the first part of MIS-3 and during the Holocene (Fig. 6a). This observation of higher chironomid diversity during warmer intervals

(early MIS-3, Holocene) across a large part of the last glacial cycle at Billjakh is in line with our results for the LGIT records (Fig. 3), which also showed increasing chironomid diversity with increasing temperature.

The sediment sequence of lake CF8, located in the northeastern Canadian Arctic, contains *in situ* preserved lake sediments that extend back from the present through the current interglacial (Holocene or Marine Isotope Stage (MIS)-1) to the two previous interglacials (MIS-5e and MIS-7). As such, it provides a unique record of past environmental change in the North American high Arctic (Axford et al., 2009). The MIS 7 part of the CF8 record is relatively short, likely recording the waning, relatively cool part of this interglacial, and contains only a few chironomid samples with count sums that allow the calculation of changes in taxon richness (Fig. 6b). High taxon richness is recorded during the early and middle part of the MIS-5e record, when temperatures inferred from chironomid assemblages are significantly higher than today (Axford et al., 2009). Rarefaction results show increases from low values in the earliest part of the Holocene to maximum values in the cold late Holocene (Axford et al., 2009) with taxon richness values averaging around 10, similar to the maximum values reached during MIS-5e. The relatively low alpha diversity across the early and middle Holocene, when past temperatures likely were highest (e.g. Gajewski, 2015; Briner et al., 2016), is likely explained by the lower taxonomic resolution of this record.

Quantitative comparison of modern-day and fossil trends in alpha diversity

This study provides, for the first time, a quantitative comparison between centennial- to multi-millennial-scale records and modern-day patterns in chironomid diversity. July air temperature shows a strong positive association to alpha diversity in our modern datasets (Fig. 2; Fig. 7), with particularly strong increases in diversity observed toward the colder end of the temperature gradient (ca. 2.5-14 °C). Similarly, most of the LGIT records presented in

this study document trends in chironomid alpha diversity that follow regional temperature variability, with higher diversity during the relatively warm Bølling/Allerød interstadial and during the early Holocene, and lower diversity during the YD and Older Dryas (Fig. 3). The ΔRF values for the modern Norwegian data and the fossil sites are mostly of a similar magnitude and show that there is a general trend of increasing ΔRF with increasing ΔT (Fig. 7a). The increase in ΔRF is limited when ΔT is low (e.g. $<2\text{ }^{\circ}\text{C}$). Between-site variation is high for the LGIT records and there is a relatively high number of fossil sites that show a ΔRF that is relatively small when compared to the ΔRF observed in the Norwegian dataset for similar ΔT and T_{jul} conditions. This low increase in ΔRF for the fossil sites relative to the ΔRF observed in the modern sites might be related to the fact that in the modern dataset there are a number of environmental factors that differ between the individual sites (e.g. geology, soil development, microclimate), whereas the variations in environmental factors with time at one particular fossil site are likely to be lower, and mainly limited to climate change and its direct effects on catchment conditions.

The results for the YD-Holocene transition also show a general pattern of increasing ΔRF with increasing ΔT in the fossil data (Fig. 7b), resembling both the pattern and magnitude of change in RF with increasing ΔT as observed in the modern Norwegian dataset as well as the pattern observed for the B/A-YD transition. Results indicate that for this transition, which is characterised by abrupt climate warming, even small-scale warming (e.g. $<2\text{ }^{\circ}\text{C}$) led to a relatively high increase in ΔRF . There are a few sites that show a ΔRF in the fossil data that is larger than ΔRF observed in the modern data (for similar ΔT and T_{jul} conditions), but again between-site variation is relatively large with some sites showing hardly any increase in ΔRF even when e.g. ΔT is $4\text{ }^{\circ}\text{C}$ or higher. Concluding, the quantitative comparison between the Norwegian dataset and the LGIT records shows that the observed changes in alpha diversity are of similar direction and magnitude (Fig. 7). They furthermore

highlight that the positive association between alpha diversity and T_{jul} is especially well-expressed in colder or more Arctic sites, and, as expected, for larger changes in temperature (e.g. $\Delta T > 4^\circ C$).

Discussion

Palaeoecological evidence for spatiotemporal trends in biodiversity

Our LGIT records generally show decreasing alpha diversity during colder periods such as the YD (Fig. 3). This confirms earlier, less comprehensive evidence on the relationship between temperature and lacustrine chironomid diversity patterns. Levesque et al. (1995) reconstructed chironomid diversity trends across the LGIT for four eastern Canadian records. Despite the low taxonomic resolution relative to later reconstructions, Levesque et al. (1995) documented decreasing alpha diversity during periods characterised by cold climate, similar to the results obtained in this study. Massaferro & Corley (1998) also reported decreasing chironomid diversity in response to climate cooling during the LGIT interval for sites in northern Patagonia. The amplitude of change in diversity in the records by Massaferro & Corley (1998) is relatively muted, which could be the result of the lower temperature amplitude associated with LGIT climate change in Northern Patagonia compared to the regions adjacent to the North Atlantic Ocean (Levesque et al., 1995; this study). However, there are major differences in the chironomid fauna adapted to cold environments in the southern hemisphere compared to the northern hemisphere (e.g. Brundin 1966), and Eggermont & Heiri (2012) show that globally different chironomid subfamilies have varying thermal optima. As chironomid-environment calibration datasets, as well as fossil datasets across a range of timescales, exist and are being developed for e.g. Australia (Rees et al., 2008), New Zealand (Van den Bos et al., 2018) or South America (Matthews-Bird et al., 2016), there is potential to apply our approach of comparing modern and fossil trends in

diversity to regions of the Southern Hemisphere as well as to the tropical regions presently under-represented in respect to fossil chironomid datasets. This will further improve our understanding of global drivers of chironomid diversity and will help to better predict future changes under projected global warming.

Stivrins et al. (2016) analysed lateglacial turnover rates in the Baltic region for a range of organisms, including higher plants, phytoplankton and animal communities. In general, all examined botanical and zoological groups show a similar trend, with lower turnover associated with the B/A-YD transition, and higher turnover associated with the YD-Holocene transition (Stivrins et al., 2016). This partially resembles our results, as we also observe abrupt and distinct changes in compositional turnover associated with Younger Dryas cooling and Holocene warming. We hypothesise that the relatively smaller impact of the cooling associated with the onset of the Younger Dryas on chironomid diversity, relative to the impact of the warming after the event, is related to the more gradual nature of this transition, whereas the onset of the Holocene in Europe was more abrupt (Brauer et al., 1999; Engels et al., 2016) and therefore had a stronger impact on freshwater ecosystems, including the chironomid fauna of our study sites.

The amplitude of temperature change during the Holocene was relatively small and occurred gradually at many of our sites, and we suggest that long term variations in site-specific factors such as soil development, vegetation change, nutrient availability and changes in the macrophyte community also influenced the development of the chironomid diversity in individual lakes. It could be argued that in the North Atlantic region the early Holocene (Walker et al., 2012) is the period with the highest environmental dynamics, as the catchments surrounding many sites were colonised by trees, soils stabilised and weathering dynamics and nutrient availability changed accordingly, potentially affecting the chironomid fauna at each of our study sites. However, our chironomid diversity records generally lack an

initial phase of instability during the early Holocene. Recent publications suggest that post-glacial ecosystem development in lakes has been much faster than previously thought (Helmens et al., 2018), and, combined with the potential for fast dispersal and colonisation by chironomids (Brooks et al., 2007), this might explain the relatively stability of our chironomid diversity records during the early Holocene.

Our results show a general trend of increasing chironomid diversity as a result of recent climate warming in two out of our three records (Fig. 5), which is in line with centennial-scale compositional turnover reconstructions for seven out of eight chironomid records from the Arctic (Smol et al., 2005). Ilyashuk et al. (2015) reconstructed trends in chironomid compositional turnover in three remote lakes in the Alps. While they show different trends for most of the past two millennia, major assemblage turnover was visible in all three sites in the interval between 1850 CE and the present (Ilyashuk et al., 2015). Finally, Nevalainen et al. (2015) studied chironomid diversity trends in four climate-sensitive treeline lakes in Austria. While their results show inter-site variability, chironomid diversity did increase in two of the studied lakes toward the twentieth and twenty-first centuries.

Effects of projected global warming on chironomid diversity

Summer temperatures are predicted to increase across the 21st century as a result of enhanced atmospheric greenhouse gas concentrations, where polar amplification will result in higher temperature increases in high-latitude regions compared to the global average (Intergovernmental Panel on Climate Change, 2014). We speculate that cold-stenothermic taxa will go locally extinct from many shallow Arctic and subarctic lakes under sustained global warming (Armitage et al., 1995; Hamerlik et al., 2017). However, our results suggest that, as a result of this projected warming, lakes in the Arctic and subarctic are likely to experience an increase in chironomid diversity, as warm-stenothermic taxa disperse into

regions and habitats that currently experience temperatures that are too low for their development.

Our palaeoecological records show that for periods with relatively small temperature change, e.g. in the order of 2 °C or less, the response of chironomid diversity to temperature change is muted or even non-existent, and the influence of factors-other-than-temperature is relatively more important. Additionally, our results show stabilising or even decreasing alpha diversity with high summer temperatures (ca. >14 °C) which could at least partially be the result of indirect climate effects, where e.g. sufficiently high temperatures could result in algal blooms, longer growing seasons, lake water stratification and decreases in hypolimnetic oxygen availability, including greater winter under-ice oxygen depletion following warmer more productive summers, favouring chironomid taxa adapted to these environmental conditions. Combined, this means that for mid-latitude regions in the Northern Hemisphere, where projected warming is relatively small compared to high-latitude regions, a temperature-driven increase in chironomid diversity across the 21st century should not necessarily be expected.

The decreasing trend in chironomid diversity at higher temperatures as seen in some of our modern datasets could furthermore be explained by the greater human impact in catchments of warmer lakes, especially in Europe, where population density and human impact on the landscape are greater in the warmer parts of the regions covered by the respective datasets. Increased nutrient influx due to human land use has been known to lead to the disappearance of aquatic macrophytes (e.g. Bradshaw et al., 2005; Engels et al., 2018), as well as to a decrease in oxygen availability (Quinlan and Smol, 2001), thus decreasing habitat availability and variability, and, ultimately, the potential for a diverse chironomid community to develop or persist. With the ongoing anthropogenic impact on the landscape (Steffen et al., 2015) we will likely see further disturbances in freshwater ecosystems across

the globe (Doncaster et al., 2016). Our results imply expected overall chironomid diversity increases under sustained global warming, but with complex and not necessarily predictable responses for individual sites due to the effects of local disturbances such as eutrophication or land use change.

Chironomids are often among the most abundant invertebrates in a lake ecosystem and are a keystone taxon in freshwater ecosystems (Brooks et al., 2007). Their preservation in lake sediment records provides one of the very few means to reconstruct changes in insect diversity on timescales longer than e.g. the last 100 years. Recent studies that highlighted the ongoing declines in insect abundance and diversity were focussed on grasslands, heathlands and wastelands and did not include aquatic groups (e.g. Hallmann et al., 2017; Jourdan et al., 2018). As freshwater communities are threatened by high rates of biodiversity loss due to a range of different environmental factors (Dudgeon et al., 2006), a better understanding of natural variability and resilience in freshwater ecosystems is critical for effective wetland management and the conservation of freshwater biodiversity. Our results illustrate the potential of using chironomids as indicators of spatiotemporal changes in biodiversity and highlight the unique perspective that this source of data can provide in order to test ecological theory about diversity response to environmental change.

Acknowledgements

We would like to thank John Birks, Konrad Gajewski, Ulrike Herzschuh, Andy Lotter, Ludmila Pestryakova and Luidmila Syrykh who played a vital role in compiling the modern-day chironomid datasets, Emmanuel Gandouin for providing access to the Aubrac dataset, and Andrew Rees for help with the GAM analyses. We acknowledge the financial support of the York University International Collaboration Grant awarded to ASM that made it possible to organise a discussion meeting that led to this study. YA and ASM acknowledge the

support of NSF Polar Programs grant 1454734. LN was supported by Deutsche
Forschungsgemeinschaft (DFG) Project NA 760/5-1 and DI 655/9-1, and by the grant of the
Russian Science Foundation (Grant 16-17-10118). We'd like to thank three anonymous
reviewers whose constructive comments helped to improve this manuscript.

The authors have no conflicts of interest.

Supporting information

Supplementary information consists of two files. First, a Word file containing three tables
(Table S1: Modern chironomid datasets; Table S2: Lateglacial-interglacial transitions sites;
Table S3: Holocene, recent and other chironomid records) and one figure (Figure S1a:
Illustration of ΔRF calculation for Norwegian dataset; Figure S1b: Illustration of ΔRF
calculation for LGIT records), as well supporting text and the R code used to produce the
results presented here. Second, the Supplementary Information contains an XLS-file with
datasheets containing the results of this study.

Data accessibility

The data produced in this study can be found in the Supplementary information. The datafile
includes all the diversity estimates produced as well as the information needed to produce the
figures presented in this manuscript. Original chironomid count datasets can be requested
from the original authors; see Supplementary Tables S1 (calibration datasets) and S2-S3
(fossil records) for the original publication details.

References

- Andrén, E., Klimaschewski, A., Self, A. E., St. Amour, N., Andreev, A. A., Bennett, K. D.,
Conley, D. J., Edwards, T. W. D., Solovieva, N., & Hammarlund, D. (2015).
Holocene climate and environmental change in north-eastern Kamchatka (Russian Far
East). *Global and Planetary Change*, 134, 41–54. doi:10.1016/j.gloplacha.2015.02.013
- Antonsson, K., Brooks, S. J., Seppä, H., Telford, R. J., & Birks, H. J. B. (2006). Quantitative
palaeotemperature records inferred from fossil pollen and chironomid assemblages
from Lake Giltjärnen, northern central Sweden. *Journal of Quaternary Science*, 21,
831–841. doi:10.1002/jqs.1004
- Armitage, P. D., Cranston, P. S., & Pinder, L. C. V. (1995). *The Chironomidae: biology and
ecology of non-biting midges*. London: Chapman and Hall.
- Axford, Y., Briner, J. P., Francis, D. R., Miller, G. H., Walker, I. R., & Wolfe, A. P. (2009).
Chironomids record terrestrial temperature changes throughout Arctic interglacials of
the past 200,000 yr. *Geological Society of America Bulletin*, 123, 1275–1287.
doi:10.1130/B30329.1
- Bartlein, P. J., Edwards, M. E., Hostetler, S. W., Shafer, S. L., Anderson, P. M., Brubaker, L.
B., & Lozhkin, A. V. (2015). Early-Holocene warming in Beringia and its mediation
by sea-level and vegetation changes. *Climate of the Past*, 11, 873–932.
doi:10.5194/cp-11-1197-2015
- Benjamini, Y., Hochberg, Y. (1995) Controlling the false discovery rate: a practical and
powerful approach to multiple testing. *Journal of the Royal Statistical Society B*, 57,
289-300. doi:10.1111/j.2517-6161.1995.tb02031.x
- Bidau, C. J. (2018). Doomsday for insects? The alarming decline of insect populations
around the world. *Entomology, Ornithology & Herpetology*, 7, e130.
doi:10.4172/2161-0983.1000e130

- Birks, H. H., & Birks, H. J. B. (2008). Biological responses to rapid climate change at the Younger Dryas-Holocene transition at Kråkenes, Western Norway. *Holocene*, 18, 19-30. doi:10.1177/0959683607085572
- Birks, H. J. B. (2007). Estimating the amount of compositional change in late-Quaternary pollen-stratigraphical data. *Vegetation History and Archaeobotany*, 16, 197-202. doi:10.1007/s00334-006-0079-1
- Birks, H. J. B., & Line, J. M. (1992). The use of rarefaction analysis for estimating palynological richness from Quaternary pollen-analytical data. *The Holocene*, 2, 1–10. doi:10.1177/095968369200200101
- Birks, H. J. B. (2012). Analysis of stratigraphical data. In: Birks, H. J. B., Lotter, A. F., Juggins, S. et al. (eds.) *Tracking Environmental Change using Lake Sediments*, Vol. 5: Data Handling and Numerical Techniques. Dordrecht, The Netherlands: Springer.
- Birks, H. J. B., Felde, V. A., & Seddon, A. W. R. (2016a). Biodiversity trends within the Holocene. *The Holocene*, 26, 994-1001. doi:10.1177/0959683615622568
- Birks, H. J. B., Felde, V. A., Bjune, A. E., Grytnes, J.-A., Seppä, H., & Giesecke, T. (2016b). Does pollen-assemblage richness reflect floristic richness? A review of recent developments and future challenges. *Review of Palaeobotany and Palynology*, 228, 1-25. doi:10.1016/j.revpalbo.2015.12.011
- Bradshaw, E. G., Rasmussen, P., & Odgaard, B. V. (2005). Mid- to late-Holocene land-use change and lake development at Dallund Sø, Denmark: synthesis of multiproxy data, linking land and lake. *The Holocene*, 15, 1152–1162. doi:10.1191/0959683605hl885rp
- Brauer, A., Endres, C., Günter, C., Litt, T., Stebich, M., & Negendank, J. F. W. (1999). High resolution sediment and vegetation responses to Younger Dryas climate change in

varved lake sediments from Meerfelder Maar, Germany. *Quaternary Science Reviews*, 18, 321-329. doi:10.1016/S0277-3791(98)00084-5

Briner, J. P., McKay, N. P., Axford, Y., Bennike, O., Bradley, R. S., de Vernal, A., Fisher, D., Francus, P., Fréchette, B., Gajewski, K., Jennings, A., Kaufmann, D. S., Miller, G., Rouston, C., & Wagner, B. (2016). Holocene climate change in Arctic Canada and Greenland. *Quaternary Science Reviews*, 147, 340-364. doi: 10.1016/j.quascirev.2016.02.010

Brooks, S. J. (2006). Fossil midges (Diptera: Chironomidae) as palaeoclimatic indicators for the Eurasian region. *Quaternary Science Reviews*, 25, 1894–1910. doi:10.1016/j.quascirev.2005.03.021

Brooks, S. J., & Birks, H. J. B. (2001). Chironomid-inferred air temperatures from Lateglacial and Holocene sites in north-west Europe: progress and problems. *Quaternary Science Reviews*, 20, 1723-1741. doi:10.1016/S0277-3791(01)00038-5

Brooks, S. J., Langdon, P. G., & Heiri, O. (2007). The Identification and Use of Palaeartic Chironomidae Larvae in Palaeoecology. *Quaternary Research Association Technical Guide no. 10*. Cambridge, UK: Quaternary Research Association.

Brooks, S. J., Davies, K. I., Mather, K. A., Matthews, I. P., & Lowe, J. J. (2016). Chironomid-inferred summer temperatures for the Last Glacial–Interglacial Transition from a lake sediment sequence in Muir Park Reservoir, west-central Scotland. *Journal of Quaternary Science*, 31, 214-224. doi:10.1002/jqs.2860

Brundin, L. (1966). Transantarctic relationships and their significance, as evidenced by chironomid midges; with a monograph of the subfamilies Podonominae and Aphroteniinae and the austral Heptagytiae. *Kungliga Svenska Vetenskapsakademiens Handlingar*, 4, 11. Stockholm: Almqvist & Wiksell.

707 Cardinale, B. J., Gonzalez, A., Allington, G. R., & Loreau, M. (2018). Is local biodiversity
708 declining or not? A summary of the debate over analysis of species richness time
709 trends. *Biological Conservation*, 219, 175-183. doi:10.1016/j.biocon.2017.12.021

710 Clegg, B. F., Kelly, R., Clarke, G. H., Walker, I. R., & Hu, F. S. (2011). Nonlinear response
711 of summer temperature to Holocene insolation forcing in Alaska. *Proceedings of the*
712 *National Academy of Sciences*, 108, 19299-19304. doi:10.1073/pnas.1110913108

713 Colombaroli, D., & Tinner, W. (2013). Determining the long-term changes in biodiversity
714 and provisioning services along a transect from Central Europe to the Mediterranean.
715 *The Holocene*, 23, 1625-1634. doi:10.1177/0959683613496290

716 Corlett, R. T. (2015). The Anthropocene concept in ecology and conservation. *Trends in*
717 *Ecology and Evolution*, 30, 36-41. doi:10.1016/j.tree.2014.10.007

718 De Vos, J. M., Joppa, L. N., Gittleman, J. L., Stephens, P. R., & Pimm, S. L. (2015).
719 Estimating the normal background rate of species extinction. *Conservation Biology*,
720 29, 452-462. doi:10.1111/cobi.12380

721 Dickson, T. R., Bos, D. G., Pellatt, M. G., & Walker, I. R. (2014). A midge-salinity transfer
722 function for inferring sea level change and landscape evolution in the Hudson Bay
723 Lowlands, Manitoba, Canada. *Journal of Paleolimnology*, 51, 325-341.
724 doi:10.1007/s10933-013-9714-x

725 Diekmann, B., Pestryakova, L., Nazarova, L., Subetto, D., Tarasov, P., Stauch, G., Thiemann,
726 A., Lehmkuhl, F., Biskaborn, B., Kuhn, G., Henning, D., & Müller, S. (2016). Late
727 Quaternary Lake Dynamics in the Verkhoyansk Mountains of Eastern Siberia:
728 Implications for Climate and Glaciation History. *Polarforschung*, 86, 97–110.
729 doi:10.2312/polarforschung.86.2.97

730 Doncaster, C. P., Alonso Chávez, V., Viguier, C., Wang, R., Zhang, E., Dong, X., Dearing, J.
731 A., Langdon, P. G., & Dyke, J. G. (2016). Early warning of critical transitions in

biodiversity from compositional disorder. *Ecology*, 97, 3079-3090.
doi:10.1002/ecy.1558

Dudgeon, D., Arthinton, A. H., Gessner, M. O., Kawabata, Z.-I., Knowler, D. J., Lévêque, C.,
Naiman, R. J., Prieur-Richard, A.-H., Soto, D., Stiasny, S. L. J., & Sullivan, C. A.
(2006). Freshwater biodiversity: importance, threats, status and conservation
challenges. *Biological Reviews*, 81, 163-182. doi:10.1017/S1464793105006950

Eggermont, H., & Heiri, O. (2012). The chironomid-temperature relationship: expression in
nature and palaeoenvironmental implications. *Biological Reviews*, 87, 430-456.
doi:10.1111/j.1469-185X.2011.00206.x

Engels, S., & Cwynar, L. C. (2011). Changes in fossil chironomid remains along a depth
gradient: evidence for common faunal breakpoints within lakes. *Hydrobiologia*, 665,
15-38. doi:10.1007/s10750-011-0601-z

Engels, S., Helmens, K. F., Välranta, M., Brooks, S. J., & Birks, H. J. B. (2010). Early
Weichselian (MIS 5d and 5c) temperatures and environmental changes in northern
Fennoscandia as recorded by chironomids and macroremains at Sokli, northeast
Finland. *Boreas*, 39, 689-704. doi:10.1111/j.1502-3885.2010.00163.x

Engels, S., Cwynar, L. C., Shuman, B. N., & Rees, A. B. H. (2012). Chironomid-based water
depth reconstructions: an independent evaluation of site-specific and local inference
models. *Journal of Paleolimnology*, 48, 693-709. doi:10.1007/s10933-012-9638-x

Engels, S., Brauer, A., Buddelmeijer, N., Martín-Puertas, C., Rach, O., Sachse, D., & van
Geel, B. (2016). Subdecadal-scale vegetation responses to a previously unknown late-
Allerød climate fluctuation and Younger Dryas cooling at Lake Meerfelder Maar
(Germany). *Journal of Quaternary Science*, 31, 741-752. doi:10.1002/jqs.2900

Engels, S., van Oostrom, R., Cherli, C., Dungait, J. A. J., Jansen, B., van Aken, J. M., van
Geel, B., & Visser, P. M. (2018). Natural and anthropogenic forcing of Holocene lake

ecosystem development at Lake Uddelermeer (The Netherlands). *Journal of Paleolimnology*, 59, 329-347. doi:10.1007/s10933-017-0012-x

Feurdean, A., Parr, C.L., Tantau, I., Farcas, S., Marinova, E., & Persoiu, I. (2013). Biodiversity variability across elevations in the Carpathians: parallel change with landscape openness and land use. *The Holocene*, 23, 869–881. doi:10.1177/0959683612474482

Fortin, M.-C., & Gajewski, K. (2011). Modern chironomid assemblages from the Canadian Arctic. *GeoHydro conference abstract*, Quebec, Canada.

Gajewski, K. (2015). Quantitative reconstruction of Holocene temperatures across the Canadian Arctic and Greenland. *Global and Planetary Change*, 128, 14-23. doi:10.1016/j.gloplacha.2015.02.003

Gajewski, K., Bouchard, G., Wilson, S. E., Kurek, J., & Cwynar, L. C. (2005). Distribution of Chironomidae (Insecta: Diptera) Head Capsules in Recent Sediments of Canadian Arctic Lakes. *Hydrobiologia*, 549, 131-143. doi:10.1007/s10750-005-5444-z

García, L. V. (2004). Escaping the Bonferroni iron claw in ecological studies. *Oikos*, 105, 657-663. doi:10.1111/j.0030-1299.2004.13046.x

Gregory-Eaves, I., & Beisner, B. E. (2011). Palaeolimnological insights for biodiversity science: an emerging field. *Freshwater Biology*, 56, 2653-2661. doi: 10.1111/j.1365-2427.2011.02677.x

Griffiths, K., Michelutti, N., Sugar, M., Douglas, M. S. V., & Smol, J. P. (2017). Ice-cover is the principal driver of ecological change in High Arctic lakes and ponds. *PLoS One*, 12, e0172989. doi:10.1371/journal.pone.0172989

Hallmann, C. A., Sorg, M., Jongejans, E., Siepel, H., Hofland, N., Schwan, H., Stenmans, W., Müller, A., Sumser, H., Hörren, T., Goulson, D., & de Kroon, H. (2017). More than

781 75 percent decline over 27 years in total flying insect biomass in protected areas.
782 PLOS One, 12, e0185809. doi:10.1371/journal.pone.0185809

783 Hamerlik, L., Svitok, M., Novikmec, M., Veslská, M., & Bitušik, P. (2017). Weak altitudinal
784 pattern of overall chironomid richness is a result of contrasting trends of subfamilies
785 in high-altitude ponds. *Hydrobiologia*, 793, 67-81. doi:10.1007/s10750-016-2992-3

786 Heiri, O., & Lotter, A. F. (2010). How does taxonomic resolution affect chironomid-based
787 temperature reconstruction? *Journal of Paleolimnology*, 44, 589–601.
788 doi:10.1007/s10933-010-9439-z

789 Heiri, O., Brooks, S. J., Renssen, H., Bedford, A., Hazekamp, M., Ilyashuk, B., Jeffers, E. S.,
790 Lang, B., Kirilova, E., Kuiper, S., Millet, L., Samartin, S., Toth, M., Verbruggen, F.,
791 Watson, J. E., van Asch, N., Lammertsma, E., Amon, L., Birks, H. H., Birks, H. J. B.,
792 Mortenson, M. F., Hoek, W. Z., Magyari, E., Munoz Sobrino, C., Seppä, H., Tinner,
793 W., Tonkov, S., Veski, S., & Lotter, A. F. (2014). Validation of climate model-
794 inferred regional temperature change for late-glacial Europe. *Nature Communications*,
795 5, 4914. doi:10.1038/ncomms5914

796 Helmens, K. F., Katrantsiotis, C., Salonen, J. S., Shala, S., Bos, J. A. A., Engels, S.,
797 Kuosmanen, N., Luoto, T. P., Väliänta, M., Luoto, M., Ojala, A., Risberg, J., &
798 Weckström, J. (2018). Warm summers and rich biotic communities during N-
799 Hemisphere deglaciation. *Global and Planetary Change*, 167, 61-73.
800 doi:10.1016/j.gloplacha.2018.05.004

801 Hooper, D. U., Chapin III, F. S., Ewel, J. L., Hector, A., Inchausti, P., Lavorel, S., Lawton, J.
802 H., Lodge, D. M., Loreau, M., Naeem, S., Schmid, B., Setälä, H., Symstad, A. J., &
803 Vandermeer, J., & Wardle D. A. (2005). Effects of biodiversity on ecosystem
804 functioning: a consensus of current knowledge. *Ecological Monographs*, 75, 3-35.
805 doi:10.1890/04-0922

806 Hölker, F., Vanni, M. J., Kuiper, J. J., Meile, C., Grossart, H.-P., Stief, P., Adrian, R., Lorke,
 807 A., Dellwig, O., Brand, A., Hupfer, M., Mooij, W. M., Nützmann, & G.,
 808 Lewandowski, J. (2015). Tube-dwelling invertebrates: tiny ecosystem engineers have
 809 large effects in lake ecosystems. *Ecological Monographs*, 85, 333-351.
 810 doi:10.1890/14-1160.1

811 Ilyashuk, E. A., Ilyashuk, B. P., Tylmann, W., Koinig, K. A., & Psenner, R. (2015).
 812 Biodiversity dynamics of chironomid midges in high-altitude lakes of the Alps over
 813 the past two millennia. *Insect Conservation and Diversity*, 8, 547–561.
 814 doi:10.1111/icad.12137

815 Intergovernmental Panel on Climate Change. (2014). *Climate Change 2014: Synthesis*
 816 *Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report*
 817 *of the Intergovernmental Panel on Climate (Geneva, Switzerland).*

818 Jourdan, J., O'Hara, R. B., Bottarin, R., Huttunen, K.-L., Kuemmerlen, M., Monteith, D.,
 819 Muotka, T., Ozoliņš, D., Paavola, R., Pilotto, F., Springe, G., Skuja, A., Sundermann,
 820 A., Tonkin, J. D., & Haase, P. (2018). Effects of changing climate on European
 821 stream invertebrate communities: A long-term data analysis. *Science of the Total*
 822 *Environment*, 621, 588-599. doi:10.1016/j.scitotenv.2017.11.242

823 Jost, L. (2007). Partitioning diversity into independent alpha and beta components. *Ecology*,
 824 88, 2427–2439. doi:10.1890/06-1736.1

825 Kaufman, D. S., Ager, T. A., Anderson, N. J., Anderson, P. M., Andrews, J. T., Bartlein, P.
 826 J., Brubaker, L. B., Coats, L. L., Cwynar, L. C., Duvall, M. L., Dyke, A. S., Edwards,
 827 M. E., Eisner, W. E., Gajewski, K., Geirsdottir, A. H. F. S., Hu, F. S., Jennings, A. E.,
 828 Kaplan, M. R., Kerwin, M. W., Lozhkin, A. V., MacDonald, G. M., Miller, G. H.,
 829 Mock, C. J., Oswald, W. W., Otto-Bliesner, B. L., Porinchu, D. F., Rühland, K.,
 830 Smol, J. P., Steig, E. J., & B Wolfe, B. B. (2004). Holocene thermal maximum in the

831 western Arctic (0–180 W). *Quaternary Science Reviews*, 23, 529-560.
832 doi:10.1016/j.quascirev.2003.09.007

833 Leather, S. R. (2018). “Ecological Armageddon”– more evidence for the drastic decline in
834 insect numbers. *Annals of Applied Biology*, 172, 1-3. doi:10.1111/aab.12410

835 Levesque, A. J., Cwynar, L. C., & Walker, I. R. (1995). Richness, diversity and succession of
836 late-glacial chironomid assemblages in New Brunswick, Canada. *Journal of*
837 *Paleolimnology*, 16, 257-274. doi:10.1007/BF00207574

838 Litt, T., Brauer, A., Goslar, T., Merkt, J., Balaga, K., Müller, H., Ralska-Jasiewiczowa, M.,
839 Stebich, M., & Negendank, J. F. W. (2001). Correlation and synchronisation of
840 Lateglacial continental sequences in northern central Europe based on annually
841 laminated lacustrine sediments. *Quaternary Science Reviews*, 20, 1233-1249.
842 doi:10.1016/S0277-3791(00)00149-9

843 Lotter, A. F., Eicher, U., Siegenthaler, U., & Birks, H. J. B. (1992). Late-glacial climate
844 oscillations as recorded in Swiss lake sediments. *Journal of Quaternary Science*, 7,
845 187-204. doi:10.1002/jqs.3390070302

846 Lotter, A. F., Birks, H. J. B., Hofmann, W., & Marchetto, A. (1997). Modern diatom,
847 cladocera, chironomid, and chrysophyte cyst assemblages as quantitative indicators
848 for the reconstruction of past environmental conditions in the Alps. I. Climate. *Journal*
849 *of Paleolimnology*, 18, 395-420. doi:10.1023/A:1007982008956

850 Lotter, A. F., Birks, H. J. B., Hofmann, W., & Marchetto, A. (1998). Modern diatom,
851 cladocera, chironomid, and chrysophyte cyst assemblages as quantitative indicators
852 for the reconstruction of past environmental conditions in the Alps. II. Nutrients.
853 *Journal of Paleolimnology*, 19, 443-463. doi:10.1023/A:1007982008956

854 Loreau, M., Naeem, S., Inchausti, P., Bengtsson, J., Grime, J. P., Hector, A., Hooper, D. U.,
855 Huston, M. A., Raffaelli, D., Schmid, B., Tilman, D., & Wardle, D. A. (2001).

Biodiversity and Ecosystem Functioning: Current Knowledge and Future Challenges.
 Science, 294, 804-808. doi:10.1126/science.1064088

Luoto, T. P. (2009a). Subfossil Chironomidae (Insecta: Diptera) along a latitudinal gradient
 in Finland: development of a new temperature inference model. *Journal of Quaternary
 Science*, 24, 150-158. doi:10.1002/jqs.1191

Luoto, T. P. (2009b). A Finnish chironomid- and chaoborid-based inference model for
 reconstructing past lake levels. *Quaternary Science Reviews*, 28, 1481-1489.
 doi:10.1016/j.quascirev.2009.01.015

Marcon, E., Hérault, B., Baraloto, C., & Lang, G. (2012). The decomposition of Shannon's
 entropy and a confidence interval for beta diversity. *Oikos*, 121, 516–522. doi:
 10.1111/j.1600-0706.2011.19267.x

Massaferro, J., & Corley, J. (1998). Environmental disturbance and chironomid
 palaeodiversity: 15 kyr BP of history at Lake Mascardi, Patagonia, Argentina. *Aquatic
 conservation: marine and freshwater ecosystems* 8, 315-323.

Matthews-Bird, F., Brooks, S. J., Holden, P. B., Montoya, E., & Gosling, W. D (2016)
 Inferring late-Holocene climate in the Ecuadorian Andes using a chironomid-based
 temperature inference model. *Climate of the Past*, 12, 1263-1280. doi:10.5194/cp-12-
 1263-2016

McGill, B. J., Dornelas, M., Gotelli, N. J., & Magurran, A. E. (2015). Fifteen forms of
 biodiversity trend in the Anthropocene. *Trends in Ecology & Evolution*, 30, 104–113.
 doi:10.1016/j.tree.2014.11.006

Medeiros, A. S., & Quinlan, R. (2011). The distribution of the Chironomidae (Insecta:
 Diptera) along multiple environmental gradients in lakes and ponds of the eastern
 Canadian Arctic. *Canadian Journal of Fisheries and Aquatic Sciences*, 68, 1511-1527.
 doi: 10.1139/f2011-076

881 Medeiros, A. S., Friel, C. E., Finkelstein, S. A., & Quinlan, R. (2012). A high resolution
 882 multi-proxy record of pronounced recent environmental change at Baker Lake,
 883 Nunavut. *Journal of Paleolimnology*, 47, 661-676. doi:10.1007/s10933-012-9589-2
 884 Medeiros, A. S., Taylor, D. J., Couse, M., Quinlan, R., Hall, R. I., & Wolfe, B. B. (2014).
 885 Biologic and nutrient responses to catchment disturbance and warming in small lakes
 886 near the Alaskan tundra-taiga boundary. *The Holocene*, 24, 1308-1319.
 887 doi:10.1177/0959683614540955
 888 Medeiros, A. S., Gajewski, K., Porinchu, D. F., Vermaire, J. C., & Wolfe, B. B. (2015).
 889 Detecting the influence of secondary environmental gradients on chironomid-inferred
 890 paleotemperature reconstructions in northern North America. *Quaternary Science*
 891 *Reviews*, 124, 265-274. doi:10.1016/j.quascirev.2015.07.010
 892 Nazarova, L., Herzsuh, U., Wetterich, S., Kumke, Th., & Pestjakova, L. (2011).
 893 Chironomid-based inference models for estimating mean July air temperature and
 894 water depth from lakes in Yakutia, northeastern Russia. *Journal of Paleolimnology*,
 895 45, 57–71. doi : 10.1007/s10933-010-9479-4
 896 Nazarova, L., Self, A. E., Brooks, S. J., van Hardenbroek, M., Herzsuh, U., & Diekmann,
 897 B. (2015). Northern Russian chironomid-based modern summer temperature data set
 898 and inference models. *Global and Planetary Change*, 134, 10-25.
 899 doi:10.1016/j.gloplacha.2014.11.015
 900 Nazarova L., Bleibtreu A., Hoff U., Dirksen V., & Diekmann B. (2017). Changes in
 901 temperature and water depth of a small mountain lake during the past 3000 years in
 902 Central Kamchatka reflected by chironomid record. *Quaternary International*, 447, 46-
 903 58 . doi: 10.1016/j.quaint.2016.10.008

904 Nevalainen, L., Luoto, T.P., Manca, M., & Weisse, T. (2015). A paleolimnological
 905 perspective on aquatic biodiversity in Austrian mountain lakes. *Aquatic Sciences*, 77,
 906 59–69. doi: 10.1007/s00027-014-0363-6

907 Oksanen, J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin, P.
 908 R., O'Hara, R. B., Simpson, G. L., Solymos, P., Stevens, M. H. H., Szoecs, E., &
 909 Wagner, H. (2015). VEGAN: Community Ecology Package (R package version 2.2–
 910 1).

911 Porinchu, D. F., Rolland, N., & Moser, K. A. (2009). Development of a chironomid-based air
 912 temperature inference model for the Central Canadian Arctic. *Journal of*
 913 *Paleolimnology*, 41, 349–368. doi:10.1007/s10933-008-9233-3

914 Porinchu, D. F. MacDonald, G. M., Moser, K. A., Rolland, N., Kremenetski, K., Seppä, H.,
 915 & Rühland, K. (2019). Evidence of abrupt climate change at 9.3 ka and 8.2 ka in the
 916 central Canadian Arctic: Connection to the North Atlantic and Atlantic Meridional
 917 Overturning Circulation. *Quaternary Science Reviews*, 219, 204-217.
 918 doi:10.1016/j.quascirev.2019.07.024

919 Primack, R. B., Miller-Rushing, A. J., Corlett, R. T., Devictor, V., Johns, D. M., Loyola, R.,
 920 Hass, B., Pakeman, R. J., & Pejchar, L. (2018). Biodiversity gains? The debate on
 921 changes in local- vs global-scale species richness. *Biological Conservation*, 219, A1-
 922 A3.

923 Quinlan, R., & Smol, J. P. (2001). Chironomid-based inference models for estimating end-of-
 924 summer hypolimnetic oxygen from south-central Ontario shield lakes. *Freshwater*
 925 *Biology*, 46, 1529-1551. doi:10.1046/j.1365-2427.2001.00763.x

926 Rada, S., Schweiger, O., Harpke, A., Kühn, E., Kuras, T., Settele, J., & Musche, M. (2019).
 927 Protected areas do not mitigate biodiversity declines: A case study on butterflies.
 928 *Diversity and Distributions*, 25, 217-224. doi:10.1111/ddi.12854

929 Rees, A. B. H., Cwynar, L. C., & Cranston, P. S. (2008). Midges (Chironomidae,
 930 Ceratopogonidae, Chaoboridae) as a temperature proxy: a training set from Tasmania,
 931 Australia. *Journal of Paleolimnology*, 40, 1159-1178. doi: 10.1007/s10933-008-9222-
 932 6
 933 Ripple, W. J., Wolf, C., Newsome, T. M., Galetti, M., Alamgir, M., Crist, E., Mahmoud, M.
 934 I., & Laurance, W. F. (2017). World scientists' warning to humanity: a second notice.
 935 *Bioscience*, 67, 1026-1028. doi:10.1093/biosci/bix125
 936 Self, A. E., Brooks, S. J., Birks, H. J. B., Nazarova, L., Porinchu, D., Odland, A., Yang, H., &
 937 Jones, V. J. (2011). The distribution and abundance of chironomids in high-latitude
 938 Eurasian lakes with respect to temperature and continentality: development and
 939 application of new chironomid-based climate inference models in northern Russia.
 940 *Quaternary Science Reviews*, 30, 1122–1141. doi:10.1016/j.quascirev.2011.01.022
 941 Self, A. E., Klimaschewski, A., Solovieva, N., Jones, V. J., Andrén, E., Andreev, A. A.,
 942 Hammarlund, D., & Brooks, S. J. (2015). The relative influences of climate and
 943 volcanic activity on Holocene lake development inferred from a mountain lake in
 944 central Kamchatka. *Global and Planetary Change*, 134, 67 – 81.
 945 doi:10.1016/j.gloplacha.2015.06.012
 946 Simpson, G. (2018). Modelling palaeoecological time series using generalized additive
 947 models. *Frontiers in Ecology and Evolution*, doi:10.1101/322248
 948 Smol, J. P., Wolfe, A. P., Birks, H. J. B., Douglas, M. S. V., Jones, V. J., Korhola, A.,
 949 Pienitz, R., Rühland, K., Sorvari, S., Antoniades, D., Brooks, S. J., Fallu, M.-A.,
 950 Hughes, M., Keatley, B. E., Laing, T. E., Michelutti, N., Nazarova, L., Nyman, M.,
 951 Paterson, A. M., Perren, A. M., Quinlan, R., Rautio, M., Saulnier-Talbot, E., Siitonen,
 952 S., Solovieva, N., & Weckström, J. (2005). Climate-driven regime shifts in the

953 biological communities of arctic lakes. *Proceedings of the National Academy of*
 954 *Sciences*, 102, 4397-4402. doi:10.1073/pnas.0500245102
 955 Solovieva, N., Klimaschewski, A., Self, A. E., Jones, V. J., Andrén, E., Andreev, A. A.,
 956 Hammarlund, D., Lepskaya, E. V., & Nazarova, L. (2015). Holocene history of
 957 Lifebuoy Lake, northwest of the Kamchatka Peninsula. *Global and Planetary Change*,
 958 134, 55–66. doi:10.1016/j.gloplacha.2015.06.010
 959 Steffen, W., Richardson, K., Rockström, J., Cornell, S. E., Fetzer, I., Bennett, E. M., Biggs,
 960 R., Carpenter, S. R., de Vries, W., de Wit, C. A., Folke, C., Gerten, D., Heinke, J.,
 961 Mace, G. M., Persson, L. M., Ramanathan, V., Reyers, B., & Sörlin, S. (2015).
 962 Planetary boundaries: Guiding human development on a changing planet. *Science*,
 963 347, 1259855. doi:10.1126/science.1259855
 964 Stivrins, N., Soininen, J., Amon, L., Fontana, S. L., Gryguc, G., Heikkilä, M., Heiri, O.,
 965 Kisielienė, D., Reitalu, T., Stančikaitė, M., Veski, S., & Seppä, H. (2016). Biotic
 966 turnover rates during the Pleistocene-Holocene transition. *Quaternary Science*
 967 *Reviews*, 151, 100-110. doi:10.1016/j.quascirev.2016.09.008
 968 Taylor, D. J., Ballinger, M. J., Medeiros, A. S., & Kotov, A. A. (2016). Climate-associated
 969 tundra thaw pond formation and range expansion of boreal zooplankton predators.
 970 *Ecography*, 39, 43-53. doi:10.1111/ecog.01514
 971 Ter Braak, C. J. F., & Šmilauer, P. (2002). *CANOCO Reference Manual and CanoDRAW for*
 972 *Windows user's guide: software for canonical community ordination (v 4.5).*
 973 Microcomputer Power, Ithaca.
 974 Tuomisto, H. (2010). A consistent terminology for quantifying species diversity? Yes, it does
 975 exist. *Oecologia*, 164, 853–860. doi:10.1007/s00442-010-1812-0
 976 Van den Bos, V., Rees, A., Newnham, R., Vandergoes, M., Wilmshurst, J., & Augustinus, P.
 977 (2018). Holocene temperature, humidity and seasonality in northern New Zealand

978 linked to Southern Hemisphere summer insolation. *Quaternary Science Reviews*, 201,
979 77-88. doi:10.1016/j.quascirev.2018.10.008

980 Vellend, M., Baeten, L., Myers-Smith, I. H., Elmendorf, S. C., Beauséjour, R., Brown, C. D.,
981 De Frenne, P., Verheyen, K., & Wipf, S. (2013). Global meta-analysis reveals no net
982 change in local-scale plant biodiversity over time. *Proceedings of the National*
983 *Academy of Sciences*, 110, 19456–19459. doi:10.1073/pnas.1312779110

984 Walker, I. R., & Cwynar, L. C. (2006). Midges and palaeotemperature reconstruction—the
985 North American experience. *Quaternary Science Reviews*, 25, 1911–1925.
986 doi:10.1016/j.quascirev.2006.01.014

987 Walker, M. J. C., Berkelhammer, M., Björk, S., Cwynar, L. C., Fisher, D. A., Long, A. J.,
988 Lowe, J. J., Newnham, R. M., Rasmussen, S. O., & Weiss, H. (2012). Formal
989 subdivision of the Holocene Series/ Epoch: a Discussion Paper by a Working Group
990 of INTIMATE (Integration of ice-core, marine and terrestrial records) and the
991 Subcommission on Quaternary Stratigraphy (International Commission on
992 Stratigraphy). *Journal of Quaternary Science*, 27, 649–659. doi:10.1002/jqs.2565

993 Wazbinski, K. E., & Quinlan, R. (2013). Midge (Chironomidae, Chaoboridae,
994 Ceratopogonidae) assemblages and their relationship with biological and
995 physicochemical variables in shallow, polymictic lakes. *Freshwater Biology*, 58,
996 2464-2480. doi: 10.1111/fwb.12223

997 Willis, K. J., & Birks H. B. J. (2006). What Is Natural? The Need for a Long-Term
998 Perspective in Biodiversity Conservation. *Science*, 314, 1261-1265.
999 doi:10.1126/science.1122667

1000 Wooller, M. J., Saulnier-Talbot, E., Potter, B. A., Belmecheri, S., Bigelow, N., Choy, K.,
1001 Cwynar, L. C., Davies, K., Graham, R. W., Kurek, J., Langdon, P., Medeiros, A.,
1002 Rawcliffe, R., Wang, Y., & Williams, J. W. (2018). A new terrestrial

1003 palaeoenvironmental record from the Bering Land Bridge and context for human
1004 dispersal. Royal Society Open Science, 5, 180145. doi:10.1098/rsos.180145
1005 W. W. F. (2016) Living Planet Report 2016. Risk and resilience in a new era. Gland,
1006 Switzerland: WWF International.
1007

Tables

*Table 1. Spearman's rank-order coefficient of correlation for rarefaction values vs environmental variables for each of the individual calibration datasets. Due to the nature of designing and producing a chironomid-climate calibration dataset, not all environmental variables were measured at each site; if individual values were missing, these sites were left out of the calculation for that specific correlation coefficient. Typically, however, a large proportion of the sites (if not all) were included in the analysis. NA: Not Available; * statistically significant at $p = 0.05$, values corrected for multiple testing within a calibration dataset using False Discovery Rate (FDR) correction (Benjamini & Hochberg, 1995) following Garcia (2004)*

	Norwegian dataset	Russian dataset	Canadian dataset	Swiss dataset	Finnish dataset
T _{jul}	0.79*	0.43*	0.61*	0.69*	0.07
Depth	0.30*	0.18*	-0.21	0.17*	-0.05
pH	-0.18	0.10	-0.02	0.08	0.22
Conductivity	0.00	-0.01	-0.06	0.52*	0.32*
TOC/DOC	0.70*	NA	0.35*	0.64*	NA
TP	NA	NA	0.17*	0.42*	NA

Figure captions

Fig. 1: Location of (a) the lakes included in the chironomid-climate calibration datasets (blue squares) and (b) the downcore study sites (yellow circles = Lateglacial sites, blue triangles = Holocene sites, green squares = subrecent sites, orange pentagons = other sites) presented in this paper. Numbers in (b) correspond to lake numbers in Supplementary tables S2 and S3 where more details on the sites are provided.




Fig. 2: Modern-day subfossil chironomid diversity as estimated using rarefaction analysis against July air temperatures ($^{\circ}\text{C}$) for datasets from (a) Norway, (b) Russia, (c) Canada, (d) Switzerland and (e) Finland (Supplementary Table S1). Solid black lines are GAM-based trendlines for rarefaction values with uncertainty intervals in light blue.




Fig. 3: Lateglacial-interglacial transition (LGIT) chironomid diversity records from northwest Europe: (a) UK sites and (b) sites from the European mainland, arranged on a S-N transect (see Fig. 1 for location of sites). Alpha diversity estimated through rarefaction analysis (RF; red solid line); compositional turnover (CT) as determined from Detrended Canonical Correspondence Analysis (DCCA) axis 1 (black solid line) in SD units (CT; bottom x-axis) and delta-CT for adjacent samples in open black bars (Δ CT; SD units, top x-axis). Directions of CT plots differ between individual sites and are selected to enable comparison against the alpha diversity records. Dark (Younger Dryas; YD) and light shading (Bølling/Allerød (B/A) and Holocene (Hol)) indicate the Lateglacial biozones as determined for each site (see Supplementary Table S2 for more information on the sites).




Fig. 4: Chironomid diversity trends in Holocene records from (a) Kamchatka (Russia), (b) other Russian sites, (c) Norway, (d) Finland, and (e) North America. All records are plotted using the age-depth models as provided in the original publications (see Supplementary Table S3 for more details on the sites). See Fig. 3 for an explanation of the symbols.




Fig. 5: Recent chironomid diversity records from northern North America. Records are plotted using the age-depth models provided in the original publications. See Supplementary Table S3 for more details on the sites, and Fig. 3 for an explanation of the symbols. Note that the x-axis for the CT plot for T15 has been reversed to better illustrate the similarity in trends between alpha diversity and compositional turnover.




Fig. 6: Chironomid diversity records from (a) Billjakh (East Siberia, Russia) and (b) CF8 (Canada). The Billjakh record spans the period from the middle of Marine Isotope Stage (MIS) 3 to the present, whereas the CF8 record covers (parts of) the past three interglacials: MIS-7 (ca 245-186 kyr BP), MIS-5E (ca 129-116 kyr BP) and the Holocene (11.6 cal kyr BP-present). The geological periods are plotted next to the diversity records for reference. See Fig. 3 for an explanation of the symbols.





Fig. 7: Quantitative comparison of changes in alpha diversity (ΔRF) with changes in temperature (ΔT) for the Norwegian dataset (continuous field in background) and the LGIT records (circles) for (a) the Bølling/Allerød-Younger Dryas transition and (b) the Younger Dryas- Holocene transition. X-axis indicates the lowest temperature of a couplet of observations; y-axis represents the difference in absolute temperatures between the couplet of observations (ΔT). Plotted values (represented by colour scale and, for fossil records, symbol size) represent the change in alpha diversity as estimated through rarefaction analysis (ΔRF) between any couplet of observations. For instance, the figure illustrates how much, on average, RF increases in the Norwegian dataset between sites for a given lower temperature (x-axis) and sites at temperatures ΔT higher (y-axis). Similarly, the results show by how much RF decreased between the B/A and the YD (Fig. 7a; results shown as absolute values to enable comparison) and increased from the YD to the Holocene (Fig 7b) for each of the LGIT sites by plotting the average temperature of the YD (x-axis), the change in temperature across the climate transition (y-axis) and ΔRF (symbol size and colour). See



1093 *main text and Supplementary Figure S1 for more information.*

1094

1095

1096